

# Comparison Between Laparoscopic Gastric Banding and Laparoscopic Sleeve Gastrectomy

**Dr. Jassim A. Fakhro, MD**

**General surgery Dept., HGH, Hamad Medical City, QATAR**

## INTRODUCTION

Obesity is one of the most diseases that considers as a global health problem, with a prevalence of >20% among the adult population in Western countries and >30% in the USA.<sup>1,2</sup> The incidence of overweight and obesity has increased and it's been identified as an epidemic associated with an increase in the diseases related to obesity, such as coronary heart disease, type 2 diabetes, hypertension, dyslipidemia, stroke, obstructive sleep apnea, , osteoarthritis and polycystic ovary syndrome. Recent a lot of studies on cancer prevention found that increased body weight was associated with an increased death rate for all cancers combined and for cancers at multiple specific sites. Thus, high body mass index (BMI) is a risk factor for higher overall mortality.

Clinicians all over the world have long been aware of the impact of obesity on health, functioning and well-being. In search of effective solutions, morbidly obese patients are increasingly turning to bariatric surgery.

There are a lot of bariatric surgeries to a chive the desirable weight loss but the restrictive procedures are more popular as Gastric banding (GB). it is characterized by minimal invasivity, total possibility of reversibility and good weight loss at long-term. On the other hand, Sleeve gastrectomy (SG), described by Hess and Marceau since 1988 during the procedure of duodenal switch and since 1993 by Johnston in an isolated form, is a less common restrictive operation for obesity, with major invasivity and a longer learning curve than GB.

## Classification of obesity

Obesity class	BMI (Kg/M2)
Normal	18.5 – 24.9
Over weight	25 – 29.9
Morbid obesity I	30.0 – 34.9
II	35.0 – 39.9
III	Over 40

## Materials and Methods

A literature search was performed using search engine Google, HighWire Press, Springer Link The following search terms was used: morbid obesity, laparoscopic gastric banding, sleve gastrectomy, out-come, effect on BMI, complication.

## Surgical Technique

*Laparoscopic Gastric banding ( GB )* Described in 1993 by *Catona*, is a surgical option that involves placing a silicone band circumferentially around the uppermost aspect of the stomach.

The band creates a small proximal pouch that empties slowly resulting in early satiety and a decreased appetite. The band is attached to an access port that is secured to the rectus muscle and can be accessed percutaneously in the office with a needle. Injection of saline into the port results in tightening of the band. This is performed on an individual basis according to weight loss and appetite. It's the most popular in Europe while in USA it has been approved by FDA in 2001.

*Sleeve Gastrectomy* (SG) described by Hess and Marceau since 1988 during the procedure of duodenal switch and since 1993 by Johnston in an isolated form, consists of vertical gastric resection of 80% capacity with exeresis of the fundus and body of the stomach linearly from the Hiss angle to 3–4 cm from the pylorus using Endo GIA staplers, which leaves a gastric residual volume ranging from 50 ml to 200 ml However, as yet, there is no agreement on the optimum residual volume.

### **Effect of surgery on weight**

Studies showed that both SG and GB have achieved a good reduction in the excess weight (*Excess weight* is defined as the difference between the actual weight and the ideal weight for longevity) Initial success in bariatric surgery is defined as a >50% loss of excess weight, or 50% EWL. In GB %EWL was at 1 and 3 years, 41.4% and 48%, respectively. While in SG it was 57.7% at 1 year and 66% at 3 years. Patients with higher BMI may require a second-stage operation later, in order to lose the rest of their excess weight if their BMI remains >45.

### **Effect on Co-morbidities**

Both of the procedure significantly improve or cured the patient co-morbidities and those changed were related to % EWL. In SG After 12 months, 57.8% of the patients were co-morbidity-free and 31.5% presented only one comorbid condition while in GB More than 63.8% of patients with sleeping apnea improved and 46.9% of them stopped using the CPAP. Table (1) shows the effect of reducing weight on different diseases.

<b>Co-morbidities</b>	<b>Improvement in %</b>
<b>Arthritis</b>	59%
<b>Asthma</b>	82%
<b>Diabetes</b>	81%
<b>GERD</b>	74%
<b>HTN</b>	49%
<b>Hyperlipidemia</b>	32%
<b>Stress incontinence</b>	80%

### **Effect of surgery on the level of serum Ghrelin**

Ghrelin, an acylated upper gastrointestinal peptide, is the only orexigenic hormone, where circulating levels decrease with feeding and increase before meals, achieving concentrations sufficient to stimulate hunger and food intake.<sup>14</sup> This hormone is primarily produced by the enteroendocrine cells of gastric mucosa and to a lesser extent from the duodenum. The procedure of SG involves resection of the gastric fundus, the predominant part of the stomach in the

production of ghrelin, resulting in less stimulation of the hunger center. A recent study by Langer<sup>15</sup> compared the ghrelin levels in patients submitted to SG and to GB, showing in patients with SG a significant decrease in plasma ghrelin at day 1 after surgery, confirmed also after 1 and 6 months, in contrast to no change found in patients with GB. Moreover, in patients with GB, the plasma ghrelin levels 1 and 6 months after surgery appeared increased compared with the preoperative levels of the same group.

## COMPLICATIONS

Complications were reported in both procedures and percentages were reasonable. LAGB is the safest bariatric operation with a mortality of 0.2%, 30-day morbidity of 5%, and delayed complication (gastric prolapse, erosion, port-tubing disconnection) rate of 12%. It has the advantage of complete reversibility by laparoscopic explanation, preservation of anatomy, and ability to perform a SG or other malabsorptive procedures.

In SG Complications occurred in ~9% of patients included trocar-site problems such as infection, hernia and hemorrhage. Other postoperative complications include urinary tract infection and atelectasis. Some studies document a leak from the anastomosis site but of major concern and were treated conservatively.

In GB and SG mortality was less than 1%.

## CONCLUSION

Both of the procedure are safe and effective in reducing weight with more advantage for SG regarding the weight loss and effect on hunger. It's clear that GB is the best procedure for people how wants to have a reversible operation, but its not for patients how are concern about a prosthesis in their bodies. SG is superior to GB in super obese and high risk patients.

## REFERENCES

1. Nahid Hamoui, MD; Gary J. Anthonie, MD; Howard S. Kaufman, MD; Peter F. Crookes, MD Sleeve Gastrectomy in the High-Risk Patient, *Obesity Surgery* 2006, vol. 16, 1445-1449
1. Jessie H. Ahroni, PhD, ARNP<sup>1,2</sup>; Kevin F. Montgomery, MD, FACS<sup>2</sup>; Brad M. Watkins, MD, FACS<sup>2</sup> Laparoscopic Adjustable Gastric Banding: Weight Loss, Co-morbidities, Medication Usage and Quality of Life at One Year.. *Obesity Surgery* 2005, vol. 15, 641-647
1. Crystine M. Lee, Paul T. Cirangle, Gregg H. Vertical gastrectomy for morbid obesity in 216 patients: report of two-year results Jossart. *Surg Endosc* (2007) 21: 1810–1816
2. Osnat Givon-Madhala, MD; Rona Spector, MD; Nir Wasserberg, MD; Nahum Beglaibter, MD; Hagit Lustigman, BA; Michael Stein, MD; Nazik Arar, MD; Moshe

- Rubin, MD. Technical Aspects of Laparoscopic SleeveGastrectomy in 25 Morbidly Obese Patients *Obesity Surgery*2007, 17, 722-727
- 3.
  4. Gianfranco Silecchia, MD, PhD; Cristian Boru, MD; Alessandro Pecchia, MD; Mario Rizzello, M; Giovanni Casella, MD1; Frida Leonetti, MD; Nicola Basso, MD Effectiveness of Laparoscopic Sleeve Gastrectomy on Co-Morbidities in Super-Obese High-Risk Patients *Obesity Surger*2006, 16, 1138-1144
  5. O. N. Tucker & S. Szomstein & R. J. Rosenthal Indications for Sleeve Gastrectomy as a PrimaryProcedure for Weight Loss in the Morbidly Obese *J Gastrointest Surg* (2008) 12:662–667
  - 6.
  7. Fàtima Sabench Pereferrer & Mercè Hernàndez Gonzàlez & Albert Feliu Rovira & Santiago Blanco Blasco & Antonio Morandeira Rivas & Daniel del Castillo Déjardin Influence of Sleeve Gastrectomy on Several Experimental Models of Obesity: Metabolic and Hormonal Implications *OBES SURG* (2008) 18:97–108
  8. John Melissas, MD; Sofia Koukouraki, MD; John Askoxylakis, MD; Maria Stathaki, MD; Markos Daskalakis, MD; Kostas Perisinakis, PhD; Nikos Karkavitsas, MD Sleeve Gastrectomy – A Restrictive Procedure? *Obesity Surgery* 2007, 17, 57-62
  - 9.
  10. J. Vidal, MD, PhD; A. Ibarzabal, MD; J. Nicolau, MD; M. Vidov, PD; S. Delgado, MD, PhD; G. Martinez, MD, PhD; J. Balust, MD, PhD; R. Morinigo, MD; A. Lacy, MD, PhD Short-term Effects of Sleeve Gastrectomy on Type 2 Diabetes Mellitus in Severely Obese Subjects *Obesity Surgery* 2007, 17, 1069-1074
  11. Italo Braghetto, MD, FACS; Owen Korn, MD, FACS; Héctor Valladares, MD; Luís Gutiérrez, MD, FACS; Attila Csendes, MD, FACS; Anfbal Debandi, MD; JaimeCastillo, MD; Alberto Rodríguez, MD; Ana Maria Burgos, MD; Luís Brunet, MD Laparoscopic Sleeve Gastrectomy: Surgical Technique, Indications and Clinical Results *Obesity Surgery*, 17, 1442-1450
  - 12.
  13. Rudolf A. Weiner, MD, PhD; Sylvia Weiner, MD; Ingmar Pomhoff, MD; Christoph Jacobi, MD, PhD; Wojciech Makarewicz, MD; Gerhard Weigand, MD. Laparoscopic Sleeve Gastrectomy – Influence of Sleeve Size and Resected Gastric Volume *Obesity Surgery*, 17, 1297-1305
  14. Andrew A. Gumbs, MD; Michel Gagner, MD; Gregory Dakin, MD; Alfons Pomp, MD. Sleeve Gastrectomy for Morbid Obesity *Obesity Surgery*, 17, 962-969
  - 15.
  16. D. Nocca & D. Krawczykowsky & B. Bomans & P. Noël & M. C. Picot & P. M. Blanc & C. de Seguin de Hons & B. Millat & M. Gagner & L. Monnier & J. M. Fabre. A Prospective Multicenter Study of 163 Sleeve Gastrectomies: Results at 1 and 2 Years *OBES SURG* (2008) 18:560–565
  17. F. B. Langer; M. A. Reza Hoda; A. Bohdjalian; F. X. Felberbauer; J. Zacherl; E. Wenzl; K. Schindler; A. Luger; B. Ludvik; G. Prager Sleeve Gastrectomy and Gastric Banding: Effects on Plasma Ghrelin Levels *Obesity Surgery*, 15, 1024-1029
  - 18.
  19. D. Cottam, F. G. Qureshi, S. G Mattar, S. Sharma, S. Holover, G. Bonanomi, R. Ramanathan, P. Schauer Laparoscopic sleeve gastrectomy as an initial weight-loss procedure for high-risk patients with morbid obesity *Surg Endosc* (2006) 20: 859–863

20. Jacques Himpens, MD; Giovanni Dapri, MD; Guy Bernard Cadière, MD, PhD A Prospective Randomized Study Between Laparoscopic Gastric Banding and Laparoscopic Isolated Sleeve Gastrectomy: Results after 1 and 3 Years *Obesity Surgery*, 16, 1450-1456
- 21.
22. Bruno M. Balsiger & Daniel Ernst & Daniel Giachino & Ruedi Bachmann & Andreas Glaetli Prospective Evaluation and 7-Year Follow-up of Swedish Adjustable Gastric Banding in Adults with Extreme Obesity *J Gastrointest Surg* (2007) 11:1470–1477
23. Luca Busetto; Gianni Segato; Francesco De Marchi; Mirto Foletto; Maurizio De Luca; Dorina Caniato; Franco Favretti; Mario Lise; Giuliano Enzi Outcome Predictors in Morbidly Obese Recipients of an Adjustable Gastric Band *Obesity Surgery*, 12, 83-92
- 24.
25. J. M. Chevallier; F. Zinzindohoué; N. Elian; A. Cherrak; J. P. Blanche; J.L. Berta; J. J. Altman; P. H. Cugnenc Adjustable Gastric Banding in a Public University Hospital: Prospective Analysis of 400 Patients *Obesity Surgery*, 12, 93-99
26. Markus Naef, MD, MBA; Ursula Naef; Wolfgang G. Mouton, MD, PhD; Hans E. Wagner, MD Outcome and Complications after Laparoscopic Swedish Adjustable Gastric Banding: 5-Year Results of a Prospective Clinical Trial *Obesity Surgery*, 17, 195-201
- 27.
28. Thomas Lang, MD; Renward Hauser, MD; Claus Buddeberg, MD; Richard Klaghofer, PhD Impact of Gastric Banding on Eating Behavior and Weight *Obesity Surgery*, 12, 100-107

For more information please log on to <http://www.laparoscopyhospital.com>